



EX VIVO EXPANSION OF HUMAN HSC WITH GENOTOXICITY-FREE SENDAI VIRUS VECTOR TRANSIENTLY EXPRESSING HOXB4 ASSESSED BY SHEEP IN UTERO TRANSPLANTATION

Shigeo Masuda¹, Tomoyuki Abe¹⁻³, Hiroshi Ban⁴, Satoshi Hayashi⁵, Hironori Takahashi^{1,5}, Makoto Inoue⁴, Mamoru Hasegawa⁴, Yoshikazu Nagao^{2,3}, Yutaka Hanazono¹

¹Division of Regenerative Medicine, Center for Molecular Medicine, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke, Tochigi 329-0498, Japan; ²Department of Agriculture, Utsunomiya University, 443 Shimokomoriya, Mohka, Tochigi 321-4415, Japan; ³United Graduate School of Agricultural Science, Tokyo University of Agriculture and Technology, 3-8-1 Harumi-cho, Fuchu-shi, Tokyo 183-8538, Japan; ⁴DNAVEC Corporation, 6 Ohkubo, Tsukuba, Ibaraki 300-2611, Japan; ⁵Department of Obstetrics and Gynecology, National Center for Child Health and Development, 2-10-1 Ohkura, Setagaya, Tokyo 157-8535, Japan